

1. Singular Values

- (a) Suppose
- $A \in \mathbb{R}^{3 \times 2}$
- is a matrix such that
- $A^\top A$
- is given by

$$A^\top A = \begin{bmatrix} 1/\sqrt{2} & -1/\sqrt{2} \\ 1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix} \begin{bmatrix} 5 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ -1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix}. \quad (1)$$

What are the singular values of A ?**Solution:** The singular values are the square roots of the eigenvalues of $A^\top A$. The eigenvalues of $A^\top A$ are 5 and 3, since those are the diagonal entries of the diagonal matrix in the spectral decomposition.Therefore, the singular values of A are $\sqrt{5}$ and $\sqrt{3}$.

- (b) Suppose that
- $B \in \mathbb{R}^{3 \times 2}$
- has singular values 0,
- $\sqrt{2}$
- , and
- $\sqrt{7}$
- . Let
- $C = \begin{bmatrix} B & -B & 3I_3 \end{bmatrix} \in \mathbb{R}^{3 \times 7}$
- , where
- $I_3 \in \mathbb{R}^{3 \times 3}$
- is the
- 3×3
- identity matrix.
- What are the singular values of C ?**

*HINT: Consider the matrix $CC^\top \in \mathbb{R}^{3 \times 3}$.***Solution:** To find the singular values of C , we consider $CC^\top \in \mathbb{R}^{3 \times 3}$. Note that we consider this matrix rather than $C^\top C \in \mathbb{R}^{7 \times 7}$ because the former is smaller, and we get the following simplification:

$$CC^\top = \begin{bmatrix} B & -B & 3I \end{bmatrix} \begin{bmatrix} B^\top \\ -B^\top \\ 3I \end{bmatrix} = 2BB^\top + 9I. \quad (2)$$

By the shift and scale properties of eigenvalues, the eigenvalues of CC^\top are $9 + 2 \times$ the eigenvalues of BB^\top . Since the eigenvalues of BB^\top are the squared singular values of B , we know that the eigenvalues of BB^\top are 0, 2, and 7. Thus the eigenvalues of CC^\top are 9, 13 and 23. Thus the nonzero singular values of C are 3, $\sqrt{13}$, and $\sqrt{23}$.